


Pelvic Floor Muscles Guide

What are they?

Pelvic floor muscles span the bottom of the pelvis and support the pelvic organs. Your pelvic floor muscles provide support for your womb, bladder and bowel all rest on the pelvic floor muscles. If you feel a heaviness or dragging you may have a prolapse, a weakness within your pelvic floor muscles.



What causes pelvic floor dysfunction?

- Child birth
- Pregnancy
- Smoking
- Heavy lifting
- Excessive weight
- Lack of exercise
- Menopause and age related changes
- Chronic constipation

How to perform pelvic floor exercises

To perform pelvic floor exercise's sit on a firm chair and lean forward. Make sure you have emptied your bladder. Squeeze the muscles that stop you passing water; you should feel a squeeze and a lift.

Don't hold your breath when you are performing the exercises. Try to isolate the front and back set of muscles to be able to improve the strength of both. Do not push down into your vagina as you are then pushing the muscles in the wrong direction. If you feel that you are not completing them correctly contact the service, as this can lead to damage.

Pelvic floor exercises need to be completed routinely, daily up to three times per day.

It can take up to 3 – 12 months to see improvement you may notice a reduction in urinary incontinence, when you cough or sneeze.

Slow: Squeeze and hold for ___ seconds.

Pelvic floor exercises need to be repeated twice per day. Always on an empty bladder.

Pelvic floor contractions need to be completed in two sections. Fast and slow.

Fast: to squeeze and release pelvic floor muscles quickly.

At the appointment your nurse will advise you on an individual set of exercises for you to perform as each patient's pelvic floor strength are different.

There is also an app called the squeezezy app which is made by physiotherapist who specialise in women's health, which helps to guide you through pelvic floor exercises and also helps to remind you to perform them.

Setting an alarm on your phone can also help to remind you to perform them.

How to check

To check if your pelvic floor strength is improving you can try stopping mid flow once a week and monitor if you can hold this for an increasing length of time.

Some ladies chose to self-examine by inserting a clean finger into their vagina and performing a set of exercises to assess their pelvic floor strength.

Bladder and bowel health

Reduce caffeine and fizzy drinks in your diet. Keep well hydrated aim for around 2 litres of fluid per day. Monitor for constipation, as this can put pressure on pelvic floor muscles, if you are straining to bear down.

Bladder and Bowel team 01422 252086

If you have any comments about this leaflet or the service you have received you can contact :

Calderdale Bladder & Bladder Service
Beechwood Community Health Centre
60B Keighley Road
Halifax HX2 8AL

Telephone: 01422 252086

www.cht.nhs.uk

If you would like this information in another format or language contact the above.

Potřebujete-li tyto informace v jiném formátu nebo jazyce, obraťte se prosím na výše uvedené oddělení

Jeżeli są Państwo zainteresowani otrzymaniem tych informacji w innym formacie lub wersji językowej, prosimy skontaktować się z nami, korzystając z ww. danych kontaktowych

ਚ ਤੁਸੀਂ ਇਹ ਜਾਣਕਾਰੀ ਕਿਸੇ ਹੋਰ ਪ੍ਰਾਚੂਪ ਜਾਂ ਭਾਸ਼ਾ ਵਿੱਚ ਲੈਣਾ ਚਾਹੁੰਦੇ ਹੋ,
ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਉਪਰੋਕਤ ਵਿਭਾਗ ਵਿੱਚ ਸਾਡੇ ਨਾਲ ਸੰਪਰਕ ਕਰੋ।

اگر آپ کو یہ معلومات کسی اور فارمیٹ یا زبان میں درکار ہوں، تو
برائے مہربانی مندرجہ بالا شعبے میں ہم سے رابطہ کریں۔

"إذا احتجت الحصول على هذه المعلومة بشكل مغاير أو مترجمة إلى لغة مختلفة فيرجى منك الاتصال بالقسم
المذكور أعلاه"