

Medicines for Children

Oxybutynin for Daytime Urinary Symptoms

This leaflet is about the use of oxybutynin for urinary symptoms including daytime wetting, frequent passing of urine and urgency to pass urine

**This leaflet is for parents and carers about how to use this medicine in children.
Our information sometimes differs from that provided by the manufacturers, because
their information is usually aimed at adults. Please read this leaflet carefully.
Keep it somewhere safe so that you can read it again.**

Name of drug

Oxybutynin

Common Brands: Cystrin, Ditropan

Modified-release tablets: Lyrinel XL

Why is it important for my child to take this medicine?

Oxybutynin relaxes the muscles of the bladder. This will mean that the bladder can hold more urine (wee) and will help your child to control their bladder so that they do not wet themselves.

What is oxybutynin available as?

- Tablets:** 2.5mg, 2.5mg, 3mg, 5mg; these contain a small amount of lactose.
- Modified-release tablets (Lyrinel XL):** 5mg, 10mg; these contain a small amount of lactose.
- Liquid medicine:** 2.5mg in 5ml.

When should I give oxybutynin?

Oxybutynin tablets or medicine may be given **two or three times** each day. There is a special, modified-release tablet called Lyrinel XL that is given once a day.

Twice each day

- If it is to be given twice each day, this should be given once in the morning and once in the evening. Ideally, these times are 10-12 hours apart, for example some time between 7 and 8am, and between 7 and 8pm.

Three times each day

- If it is to be given three times each day, this should be first thing in the morning, early afternoon (or after school) and at bedtime. Ideally, these times should be at least 4 hours apart.

Modified-release (Lyrinel XL) tablets

- These tablets are given only **once a day**.

Give the medicine at about the same times each day so that this becomes part of your child's daily routine, which will help you to remember.

How much should I give?

Your doctor will work out the amount of oxybutynin (the dose) that is right for **your** child. The dose will be shown on the medicine label.

It is important that you follow your doctor's instructions about how much to give.

How should I give it?

Tablets should be swallowed with a glass of water, milk or juice. Your child should not chew the tablet.

Modified-release tablets (Lyrinel XL) should be swallowed with a glass of water, milk or juice. Your child should not chew the tablet, because this will change how it works.

Liquid medicine: Measure out the right amount using an oral syringe or medicine spoon. You can get these from your pharmacist. Do not use a kitchen teaspoon as it will not give the right amount.

When should the medicine start working?

It may take a few days (sometimes a week or more) for the medicine to help your child's bladder symptoms. Continue to give the medicine as your doctor has told you to during this time.

What if my child is sick (vomits)?

- If your child is sick less than 30 minutes after having a dose of oxybutynin, give them the same dose again.
- If your child is sick more than 30 minutes after having a dose of oxybutynin, you do not need to give them another dose. Wait until the next normal dose.

If your child is sick again, seek advice from your GP, pharmacist or hospital. They will decide what to do based on your child's condition and the specific medicine involved.

What if I forget to give it?

If you normally give oxybutynin twice a day

If you remember up to 4 hours after you should have given a dose, give your child the missed dose. For example, if you usually give a dose at about 7am, you can give the missed dose at any time up to 11am. If you remember after that time, do not give the missed dose. Wait until the next normal dose.

If you normally give oxybutynin three times a day

If you miss a dose, wait until the next normal dose. Do not give the missed dose.

Modified-release (Lyrinel XL)

If you forget to give a dose, you can give the missed dose first thing in the morning, as long as this is at least 12 hours before the next normal bedtime dose. If you remember after this, do not give the missed dose. You do not need to wake your child up to give them the missed dose.

Never give a double dose of oxybutynin

What if I give too much?

If you think you may have given your child too much oxybutynin, contact your doctor or local NHS services (details at end of leaflet). Have the medicine packet with you if you telephone for advice.

Are there any possible side-effects?

We use medicines to make our children better, but sometimes they have other effects that we don't want (side-effects).

Side-effects you must do something about

- Your child may sweat less whilst taking oxybutynin. Be careful that they don't overheat, particularly in hot weather and during exercise; make sure that they drink plenty of water.
- Your child may get a fast or irregular heartbeat (a fluttery feeling in the chest). Contact your doctor straight away if this happens.
- Your child's skin may become more sensitive to sunlight. Keep them out of strong sun. When outdoors they should wear long-sleeved tops and trousers and a hat, and use high-factor sun block (at least SPF 15).

Other side-effects you need to know about

- Some children get a dry mouth for the first few days. Eating citrus fruits (eg. oranges) and taking sips of water may help. This effect usually settles down quite quickly. If it doesn't contact your doctor.
- Your child may feel sick (nausea) and get diarrhoea or constipation (difficulty doing a poo). If these effects cause problems, your doctor may give you some other medicines to help.
- Your child may have some problems passing urine (doing a wee). If this happens, contact your doctor.
- Some children may feel sleepy, get a headache, have changes in mood, see strange things (hallucinations), or get nightmares (although this is rare). Some children find their eyesight is blurry or double.

These side-effects should get better after a week. If they don't contact your doctor.

There may, sometimes, be other side-effects that are not listed above. If you notice anything unusual and are concerned, contact your doctor. You can report any suspected side-effects to a UK safety scheme at <http://www.mhra.gov.uk/yellowcard>

Can other medicine be given at the same time as oxybutynin?

- You can give your child medicines that contain paracetamol or ibuprofen, unless your doctor has told you not to.
- Check with your doctor or pharmacist **before** giving any other medicines to your child. This includes herbal or complementary medicines.

General advice about medicines

- Try to give the medicine at about the same time(s) each day, to help you remember.
- Only give this medicine to your child. Never give it to anyone else, even if their condition appears to be the same, as this could do harm.
- If you think someone else may have taken the medicine by accident, contact your doctor straight away
- Make sure that you always have enough medicine. Order a new prescription at least 2 weeks before you will run out.
- Make sure that the medicine you have at home has not reached the 'best before' or 'use by' date on the packaging. Give old medicines to your pharmacist to dispose of.

Where should I keep this medicine?

- Keep the medicine in a cupboard, away from heat and direct sunlight. It does not need to be kept in the fridge.
- Make sure that children cannot see or reach the medicine.
- Keep the medicine in the container it came in.

Who to contact for more information

Your child's doctor, pharmacist or nurse will be able to give you more information about oxybutynin and about other medicines used to treat daytime urinary symptoms.

You can also get useful information from:

England: NHS 111

Tel: 111 - www.nhs.uk

Scotland: NHS 24

Tel: 111 - www.nhs24.scot

Wales: NHS 111 Wales

Tel: 111 - www.111.wales.nhs.uk

Northern Ireland: NI Direct

www.nidirect.gov.uk

ERIC (Children's Bowel & Bladder Charity)

Helpline: 0808 169 9949 www.eric.org.uk

www.medicinesforchildren.org.uk



If you have any comments about this leaflet or the service you have received you can contact :

Pharmacy Department
Calderdale Royal Hospital
Telephone: 01422 225625

www.cht.nhs.uk

If you would like this information in another format or language contact the above.

Potřebujete-li tyto informace v jiném formátu nebo jazyce,
obraťte se prosím na výše uvedené oddělení

Jeżeli są Państwo zainteresowani otrzymaniem tych
informacji w innym formacie lub wersji językowej,
prosimy skontaktować się z nami, korzystając z ww.
danych kontaktowych

ਜੇ ਤੁਸੀਂ ਇਹ ਜਾਣਕਾਰੀ ਕਿਸੇ ਹੋਰ ਪ੍ਰਾਚੂਪ ਤਾਂ ਭਾਸ਼ਾ ਵਿੱਚ ਲੈਣਾ ਚਾਹੁੰਦੇ ਹੋ,
ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਉਪਰੋਕਤ ਵਿਭਾਗ ਵਿੱਚ ਸਾਡੇ ਨਾਲ ਸੰਪਰਕ ਕਰੋ।

اگر آپ کو یہ معلومات کریں اور فارمیٹ یا زبان میں درکار ہوں، تو
برائے مہربانی مندرجہ بالا شعبے میں ہم سے رابطہ کریں۔

**"إذا احتجت الحصول على هذه المعلومة بشكل مغاير أو مترجمة إلى لغة مختلفة فيرجى منك الاتصال بالقسم
المذكور أعلاه"**