

Anaesthetic Department

You and Your Anaesthetic

Information to help patients prepare for an anaesthetic www.rcoa.ac.uk/patientinfo

This leaflet gives information to help you prepare for your anaesthetic. It has been written by patients, patient representatives and anaesthetists, working together.

You can find more information leaflets on the website www.rcoa.ac.uk/patientinfo. The leaflets may also be available from the anaesthetic department or pre-operative assessment clinic in your hospital.

The website includes the following:

- Anaesthesia explained (a more detailed booklet)
- Your spinal anaesthetic
- Anaesthetic choices for hip or knee replacement
- Epidural pain relief after surgery
- Local anaesthesia for your eye operation
- Your child's general anaesthetic
- Your anaesthetic for major surgery
- Your anaesthetic for a broken hip
- Brachial plexus block for surgery and pain relief

Risks associated with your anaesthetic

This is a collection of 14 articles about specific risks associated with having an anaesthetic or an anaesthetic procedure. It supplements the patient information leaflets listed above and is available on the website: www.rcoa.ac.uk/patients-and-relatives/risks



Throughout this leaflet and others in the series, we have used this symbol to highlight key facts.

What is anaesthesia?

Anaesthesia stops you feeling pain and other sensations. It can be given in various ways and does not always make you unconscious.

Local anaesthesia involves injections that numb a small part of your body. You stay conscious but free from pain.

Regional anaesthesia involves injections that numb a larger or deeper part of the body. You stay conscious but free from pain. For some surgery you may be aware of pressure sensations.

General anaesthesia gives a state of controlled unconsciousness. It is essential for some operations and procedures. You are unconscious and feel nothing.

Sedation gives a sleep like state and is often used with a local or regional anaesthetic. Sedation may be light or deep and you may remember everything, something or nothing after.

Anaesthetists

Anaesthetists are doctors with specialist training who:

- discuss the type or types of anaesthetic that are suitable for your operation. If there are choices available, your anaesthetist will help you choose what is best for you
- discuss the risks of anaesthesia with you
- agree a plan with you for your anaesthetic and pain control
- are responsible for giving your anaesthetic and for your wellbeing and safety throughout your surgery
- manage any blood transfusions you may need
- plan your care, if needed, in the intensive care unit
- make your experience as calm and pain free as possible.

The pre-operative assessment clinic

If you are having a planned operation (rather than emergency) you will usually be invited to a pre-operative assessment clinic a few weeks or days before your surgery.

Staff at the clinic will assess your general health. You will be asked questions. Tests will be organised if necessary. This will often include blood tests and an ECG (heart tracing).

You are likely to be checked for certain important infections – usually by using a swab on your skin, or in your nose.



Please bring a list and your medicines in their boxes.

The staff will want to make an accurate list of the medicines you take.

If you are allergic to anything, this will be written down.

If necessary, arrangements may be made for you to see an anaesthetist.

If you may need a blood transfusion during or after your operation, a blood test will be needed to prepare for this. This can only be performed a certain period/time before. You can ask for information about the risks and benefits of blood transfusion. Blood transfusions are generally avoided unless necessary.



If you know you have high blood pressure, it is a good idea to bring a list of any recent blood pressure readings. This is because blood pressure can rise in a hospital clinic, and this may not be a true picture of your usual blood pressure.

More information

This is a very useful opportunity for you to ask any questions that you have about the anaesthetic and about coming into hospital generally. If the staff do not have all the answers you need, they will be able to help you find out more.

Before coming to hospital

- If you smoke, giving up for several weeks before the operation will reduce the risk of breathing problems during your anaesthetic, making your anaesthetic safer. It is also important that you consider any mental health concerns such as anxiety and depression, as these too can make a difference to your surgery and recovery. Increasing your activity in the weeks before surgery can improve your heart function and fitness.
- If you are very overweight, reducing your weight will reduce many of the risks of having an anaesthetic.
- If you have loose teeth or crowns, treatment from your dentist may reduce the risk of damage to your teeth during the anaesthetic.
- If you have a long-standing medical problem such as diabetes, asthma or bronchitis, thyroid problems, heart problems or high blood pressure, you should ask your GP if you need a check up.

On the day of your operation



Nothing to eat or drink – ('nil by mouth')

The hospital should give you clear instructions about eating and drinking. These instructions are important. If there is food or liquid in your stomach during your anaesthetic, it could come up into your throat and damage your lungs.

If you are a smoker you should not smoke on the day of your operation.

If you are taking medicines: most medicines should be continued before an operation, but there are some important exceptions. You will need specific instructions from the pre-operative assessment team about your tablets.

If you feel unwell when you are due to come into hospital, please telephone the ward for advice.

Meeting your anaesthetist

You will meet your anaesthetist before your operation. They will refer to the discussion you had in the pre-operative assessment clinic. They will discuss the anaesthetic you could have, including benefits, risks and your preferences.

They will then:

- decide with you which anaesthetic would be best for you
- decide for you, if you would prefer that.

If there is a choice of anaesthetic, this will depend on:

- the operation you are having and your physical condition
- your preferences and the reasons for them
- the recommendation of the anaesthetist, based on experience
- the equipment, staff and resources at the hospital.

Pre-medication (a 'pre-med') is given before some anaesthetics.

Premeds prepare your body for surgery – they may start off the pain relief, reduce acid in the stomach or help you relax. Some pre-meds make you more drowsy after the operation. If you think a pre-med would help you, please ask your anaesthetist.

A needle is used to start most anaesthetics in adults. If you are very worried about this, please talk to your anaesthetist.



Sedation. This is the use of small amounts of anaesthetic or similar drugs to produce a 'sleep like' state. If you are having a local or regional anaesthetic, you will need to decide whether you would prefer to:

- be fully alert
- be relaxed and sleepy (sedation) but not unconscious have a general anaesthetic as well (if appropriate)



Nothing will happen to you until you understand and agree with what has been planned. You have the right to refuse if you do not want the treatment suggested or if you want more information or more time to decide.

When you are called for your operation

- A member of staff will go with you to the theatre.
- You can wear your glasses, hearing aids and dentures until you are in the anaesthetic room. You may be able to keep them on if you are having a local or regional anaesthetic.
- Jewellery and/or any decorative piercing should be removed, depending on your planned surgery.
- If you cannot remove it, the nurses will cover it with tape to prevent damage to it or to your skin.
- If you are having a local or regional anaesthetic, you can take your own electronic device, with headphones to listen to music.
- You may walk to theatre, accompanied by a member of staff, or you may go in a wheelchair or on a bed or trolley. If you are walking, you can wear your own dressing gown and slippers.
- **Final checks** will be done as you arrive in the operating department, before the anaesthetic starts. You will be asked to confirm your name, DOB, the operation you are having, whether left or right side (if applicable), when you last ate or drank and your allergies. These routine checks are normal in all hospitals.

Starting the anaesthetic

Your anaesthetic may start in the anaesthetic room or in the operating theatre. Your anaesthetist will be working with a trained assistant. The anaesthetist or the assistant will attach machines that measure your heart rate, blood pressure and oxygen levels.

Almost all anaesthetics, including some kinds of local anaesthetic, start with a needle being used to put a cannula (thin plastic tube) into a vein in the back of your hand or arm. If needles worry you, please tell your anaesthetist. A needle cannot usually be avoided, but there are things he/she can do to help.

Local and regional anaesthetics

If you are having a local or regional anaesthetic:

- your anaesthetist will ask you to keep quite still while the injections are given. You may notice a warm tingling feeling as the anaesthetic begins to take effect
- your operation will only go ahead when you and your anaesthetist are sure that the area is numb
- you will remain alert and aware of your surroundings, unless you are having sedation. A screen shields the operating site, so you will not see the operation unless you want to
- if you are having sedation, you will be sleepy and relaxed. However, you may be aware of events around you
- for regional anaesthetics, your anaesthetist is always near to you and you can speak to him or her whenever you want to. For local anaesthetics, other theatre staff may be looking after you

General anaesthetics

There are two ways of starting a general anaesthetic:

- anaesthetic drugs may be injected into a vein through the cannula. This is generally used for adults
- you can breathe anaesthetic gases and oxygen through a mask, which you may hold if you prefer.

Once you are unconscious, an anaesthetist stays with you at all times and continues to give you drugs to keep you anaesthetised.

The Recovery room: After the operation, you will usually be taken to the recovery room. Recovery staff will make sure you are as comfortable and free of nausea as possible. When they are satisfied that you have recovered safely from your anaesthetic you will be taken back to the ward.

Pain relief afterwards

Good pain relief is important and some people need more pain relief than others. It is much easier to relieve pain if it is dealt with before it gets bad. Pain relief can be increased, given more often, or given in different combinations.

Occasionally, pain is a warning sign that all is not well; therefore, you should always report it to your nurses and seek their advice and help.

Here are some ways of giving pain relief:

- **Pills, tablets or liquids to swallow**

These are used for all types of pain. They take at least half an hour to work. You need to be able to eat, drink and not feel sick, for these drugs to work.

- **Injections**

These are often needed, and may be intravenous (through your cannula into a vein for a quicker effect) or intramuscular (into your leg or buttock muscle using a needle, taking about 20 minutes to work).

- **Suppositories**

These waxy pellets are put in your rectum (back passage). The pellet dissolves and the drug passes into the body. They are useful if you cannot swallow or if you might vomit.

- **Patient-controlled analgesia (PCA)**

A machine allows you to control your pain relief yourself. The medicine enters your body through your cannula. If you would like more information, ask for a leaflet on PCA.

- **Local anaesthetics and regional blocks**

These types of anaesthesia can be very useful for relieving pain after surgery. More details can be found in the leaflet Epidural Pain Relief After Surgery.

Pain relief after leaving hospital

Although you may be given a supply of painkillers when you leave the hospital, it is sensible to buy some over-the-counter painkillers to have ready at home. If you are still needing them two weeks after discharge from the hospital, you should get in touch with your GP to discuss this further.

You may go home on morphine-type pain killers following your operation. It is important that you reduce and then stop these medications as soon as possible as their continued use can cause you significant harm.

Understanding risk

In modern anaesthesia, serious problems are uncommon.

Risk cannot be removed completely, but modern equipment, training and drugs have made it a much safer procedure in recent years.


To understand a risk, you must know:

- how likely it is to happen
- how serious it could be
- how it can be treated.

The risk to you as an individual will depend on:

- whether you have any other illness
- personal factors, such as smoking or being overweight
- surgery that is complicated, long or done in an emergency.

More information about risks associated with having an anaesthetic can be found on www.rcoa.ac.uk/patientinfo.



Side effects and complications

More information on the side effects and complications than is listed here is given in the booklet Anaesthesia explained (www.rcoa.ac.uk/patientinfo/risks/risk-leaflets)

RA = This may occur with a regional anaesthetic

GA = This may occur with a general anaesthetic

Very common and common side effects

RA GA	Feeling sick and vomiting after surgery
GA	Sore throat
RA GA	Dizziness, blurred vision
RA GA	Headache
RA GA	Bladder problems
GA	Damage to the lips or tongue (usually minor)
RA GA	Itching
RA GA	Aches, pains and backache
RA GA	Pain during injection of drugs
RA GA	Bruising and soreness
GA	Confusion or memory loss

Uncommon side effects and complications

GA	Chest infection
GA	Muscle pains
RA GA	Slow breathing (depressed respiration)
GA	Damage to teeth
RA GA	An existing medical condition getting worse
GA	Awareness (becoming conscious during your operation)

Rare or very rare complications

GA	Damage to the eyes
RA GA	Heart attack or stroke
RA GA	Serious allergy to drugs
RA GA	Nerve damage
RA GA	Death
RA GA	Equipment failure

Deaths caused by anaesthesia are very rare. There are probably about five deaths for every million anaesthetics in the UK.

Questions you may like to ask your anaesthetist

- 1 Who will give my anaesthetic?
- 2 Do I have to have a general anaesthetic?
- 3 What type of anaesthetic do you recommend?
- 4 Have you often used this type of anaesthetic?
- 5 Will I be unconscious and completely unaware during this kind of anaesthetic?
- 6 What are the risks of this type of anaesthetic?
- 7 Do I have any special risks?
- 8 How will I feel afterwards?

If you have any comments about this leaflet or the service you have received you can contact :

Pre-operative Assessment Department
Huddersfield Royal Infirmary

Telephone: (01484) 342527

www.cht.nhs.uk

If you would like this information in another format or language contact the above.

Potřebujete-li tyto informace v jiném formátu nebo jazyce, obraťte se prosím na výše uvedené oddělení

Jeżeli są Państwo zainteresowani otrzymaniem tych informacji w innym formacie lub wersji językowej, prosimy skontaktować się z nami, korzystając z ww. danych kontaktowych

ਬ ਤੁਸੀਂ ਇਹ ਜਾਣਕਾਰੀ ਕਿਸੇ ਹੋਰ ਪ੍ਰਾਚੂਰ ਜਾਂ ਭਾਸ਼ਾ ਵਿੱਚ ਲੈਣਾ ਚਾਹੁੰਦੇ ਹੋ,
ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਉਪਰੋਕਤ ਵਿਭਾਗ ਵਿੱਚ ਸਾਡੇ ਨਾਲ ਸੰਪਰਕ ਕਰੋ।

اگر آپ کو یہ معلومات کسی اور فارمیٹ یا زبان میں درکار ہوں، تو
برائے مہربانی مندرجہ بالا شعبے میں ہم سے رابطہ کریں۔

"إذا احتجت الحصول على هذه المعلومة بشكل مغاير أو مترجمة إلى لغة مختلفة فيرجى منك الاتصال بالقسم
المذكور أعلاه"